



Health & Wellbeing

Access to green space improves our mental wellbeing², reducing the need to treat for anxiety and mental health conditions³. Depressive disorders are now the foremost cause of disability in middle- and high-income countries⁴ and can be precursors for chronic physical health problems.

Spending time in green spaces has been shown to produce levels and patterns of chemicals in the brain associated with low stress⁵ and positive impacts on blood pressure⁶. Positive links have also been demonstrated between how well people perform at attention-demanding tasks and time spent, either beforehand or during, in green space⁶⁻⁹.

Across Europe, approximately 1 in every 15 deaths is associated with a lack of physical activity¹⁰. In the UK, only one third of the population achieves the recommended level of exercise¹¹ and the impact of this on our health is estimated to have a direct economic cost of £1 billion per year¹².

Green areas encourage physical activity by providing a pleasant environment in which to exercise¹³; linear woodland trails encourage

walking and cycling, whilst large sport and community parks encourage more formal physical activity¹⁴.

Where green space is available, the socioeconomic position of the local population does not affect how frequently it is used¹⁵, implying that where accessible green space is provided it will be used and may help to reduce socioeconomic health inequalities^{16,17}.

Urban green spaces provide pleasant areas to relax and socialise, promoting greater levels of social activity and stronger neighbourhood relationships¹⁸. This can be particularly important in maintaining a high quality of life for elderly people¹⁹⁻²¹.



Temperature & Climate Change

In the UK, urban temperatures are typically 1-2°C higher than the surrounding rural areas^{23,24}. This urban heat island (UHI) effect occurs because the materials used to build towns and cities absorb more of the sun's energy than the natural surfaces they replaced.

The UHI effect makes people living in urban areas particularly vulnerable to heat waves, for example there was an estimated 42% increase in mortality in London during the heatwave that affected Europe in August 2003²⁵.


Urban green spaces reduce the UHI effect by providing shade and by cooling the air through the process of evapotranspiration. During evapotranspiration, the sun's energy is used to transfer water from the leaves of plants into the atmosphere²⁶.

Urban green spaces are on average around 1°C cooler, during both the day and night time, than built-up regions in the same town or city²⁷, and this cooling effect can extend beyond the green space itself, into the surrounding urban areas²⁸. During the summer this may reduce the need for air conditioning, and associated energy use, in nearby buildings²⁹.

The amount of carbon dioxide in the atmosphere has increased by more than 40% since humans began industrialising, resulting in a gradual warming of the planet over the past century³⁰. Trees and plants take carbon dioxide from the atmosphere and around half of it is stored in their branches and roots, with large amounts of carbon also stored by the surrounding soils.

This process is known as carbon sequestration and, as long as the vegetation is preserved, results in an overall reduction of atmospheric carbon dioxide concentrations. However, the decomposition of dead trees and plants returns carbon dioxide to the atmosphere. Understanding the carbon balance of any green space therefore requires an analysis of the relative amounts of sequestration and decomposition, in addition to any maintenance related greenhouse gas emissions (e.g., through mowing, irrigation and the use of fertiliser).

Overall, urban green spaces take in more carbon than they return to the atmosphere^{31,32} but their design and maintenance play a crucial role in determining how much carbon they will store. For example, a "forest-like" green space with many trees and native vegetation ground cover maximises carbon sequestration over a "park-like" design with fewer trees and frequently mown grass³³. As well as creating new green space, looking after existing mature trees is particularly important because they continue to sequester and store large amounts of carbon³⁴.







Wildlife & Habitats

Our towns and cities are typically considered to host a less diverse range of plants, animals and birds than nearby rural areas⁷⁰. However, green spaces within an urban area can be home to many of the same species that are more commonly associated with rural settings⁷¹, including those that are rare or threatened^{72,73}. For some species, urban areas can provide a more favourable habitat than intensively farmed countryside^{72,74}, suggesting that towns and cities could make an important contribution to national conservation efforts.

Large parks and woodland regions are able to support the widest range of species⁷¹, but even small areas of vegetation such as roundabouts⁷⁵, roadside verges⁷⁶ and green roofs^{77,78} can support a range of plants, insects and birds.

For many city dwellers, spending time in urban green spaces is their only regular opportunity to be surrounded by nature. Research suggests that people get more enjoyment from spending time in green space when they perceive there to be a high level of biodiversity⁷⁹ and that visitors to green spaces would be willing to pay to see an enhancement in the species richness of plants, birds and invertebrates⁸⁰.

Urban green spaces can act as "wildlife corridors", linking together larger parks, and providing links to rural areas on the outskirts of towns and cities. This facilitates the movement of animals, birds and insects between individual green spaces and prevents the fragmentation and isolation of wildlife^{81,82}.

In the UK, urban green spaces form an important habitat for pollinators, such as bees, butterflies and hoverflies⁷⁴. Maintaining a healthy population of pollinators is vitally important as many flowers and crops (including tomatoes, apples and strawberries) depend upon them in order to reproduce. Pollinator populations are declining in the UK^{83,84}, so the provision of viable habitats in urban regions could form part of a broader strategy to combat this trend.

The more green space the better for urban wildlife, but strategies designed to enhance biodiversity will depend on the location, type of habitat and species present⁸⁵. However, some general themes emerge, such as: less intensive management practices, e.g., infrequent mowing of grass; protecting some parts of the green space from human interference, e.g., routing paths away from the most suitable nesting locations to prevent adverse effects on the reproductive success of birds; and the introduction of locally native wildflowers⁸⁵.

INTERCONNECTED GREEN SPACES COMPOSED OF NATIVE TREE AND SHRUB SPECIES, WITH LESS INTENSIVE MANAGEMENT, OFFER THE GREATEST BENEFITS FOR WILDLIFE





The presence of green space affects an urban region in the many different ways described in this document; the economic impacts of which are not straightforward to quantify and estimates can vary widely⁸⁶. In terms of direct financial impacts, case studies from around the UK suggest that proximity to green space is positively linked to both commercial and residential property prices, with properties overlooking a park being valued around 5-7% higher than equivalent properties elsewhere⁸⁷.

The creation, maintenance and management of green space also generates employment opportunities, and may have indirect benefits to local economies by encouraging further investment and property development in the area⁸⁶.

An assessment conducted for the Mersey Forest, a tree planting programme that now forms a 1300 km² network of woodlands and green spaces across Cheshire and Merseyside, concluded that every £1 invested in the programme was more than doubled88. This was due mostly to tourism expenditure, the creation of forestry related jobs, estimated social cost savings (such as the impact of reduced air pollution), and well-being benefits (such as people's perception of increased biodiversity and

improved visual quality of the environment). The assessment concluded that the location of green space is key; to maximise the benefits, green space must be easily accessible to both local people and tourists, or at least viewable from their homes or while travelling.

However, it is not clear whether the assignment of monetary values can fully capture the importance of non-monetary effects, such as increased biodiversity or the cultural significance of woodland. Further research is required to develop approaches that can combine both monetary and non-monetary valuations in order to assess the true value of urban green spaces.



References

- United Nations (Department of Economic and Social Affairs Population Division). World Urbanization Prospects: The 2014 Revision, Highlights (ST/ESA/ SER.A/352). (2014).
- 2. White, M. P. et al. Psychological Science, 24, 920-928, (2013).
- 3. Nutsford, D. et al. Public Health, 127, 1005-1011, (2013).
- 4. World Health Organisation. The Global Burden of Disease: 2004 update. (2008).
- Ward Thompson, C. et al. Landscape and Urban Planning, 105, 221-229. (2012).
- 6. Hartig, T. et al. Journal of Environmental Psychology, 23, 109-123, (2003).
- 7. Hartig, T. et al. Environment and Behavior, 23, 3-26, (1991).
- 8. Tennessen, C. M. & Cimprich, B. Journal of Environmental Psychology, 15, 77-85, (1995).
- 9. Roe, J. & Aspinall, P. Health & Place, 17, 103-113, (2011).
- 10. Ekelund, U. et al. The American Journal of Clinical Nutrition, (2015).
- 11. Department of Health. Start Active, Stay Active: A report on physical activity for health from the four home countries' Chief Medical Officers. (2011).
- 12. Scarborough, P. et al. Journal of Public Health, 33, 527-535, (2011).
- 13. Coombes, E. et al. Social Science & Medicine, 70, 816-822, (2010).
- 14. Brown, G. et al. Landscape and Urban Planning, 121, 34-44, (2014).
- 15. Grahn, P. & Stigsdotter, U. A. Urban Forestry & Urban Greening, 2, 1-18, (2003).
- 16. Mitchell, R. & Popham, F. The Lancet, 372, 1655-1660, (2008).
- 17. Mitchell, R. J. et al. American Journal of Preventive Medicine, (2015).
- 18. Sullivan, W. C. et al. Environment and Behavior, 36, 678-700, (2004).
- 19. Sugiyama, T. et al. Environment and Behavior, 41, 3-21, (2009).
- 20. Kweon, B.-S. et al. Environment and Behavior, 30, 832-858, (1998).
- 21. Sugiyama, T. & Ward Thompson, C. Environment and Planning A, 39, 1943-1960, (2007).
- Dallimer, M. et al. International Journal of Environmental Research and Public Health, 11, 7977-7992, (2014).
- Watkins, R. et al. Building Services Engineering Research and Technology, 23, 207-213, (2002).
- 24. Jones, P. D. & Lister, D. H. Weather, 64, 323-327, (2009).
- 25. Johnson, H. et al. Euro Surveill., 10, 558, (2005).
- Grimmond, C. S. B. & Oke, T. R. Water Resources Research, 27, 1739-1755, (1991).
- 27. Bowler, D. E. et al. Landscape and Urban Planning, 97, 147-155, (2010).
- 28. Yu, C. & Hien, W. N. Energy and Buildings, 38, 105-120, (2006).
- 29. McHale, M. R. et al. Urban Forestry & Urban Greening, 6, 49-60, (2007).
- IPCC. in Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change (eds T. F. Stocker et al.) (Cambridge University Press, Cambridge, UK and New York, NY, USA., 2013).
- 31. Nowak, D. J. et al. Environmental Pollution, 178, 229-236, (2013).
- 32. Nowak, D. J. & Crane, D. E. Environmental Pollution, 116, 381-389, (2002).
- 33. Strohbach, M. W. et al. Landscape and Urban Planning, 104, 220-229, (2012).
- 34. Stephenson, N. L. et al. Nature, 507, 90-93, (2014).
- 35. Sani, S. Energy and Buildings, 15, 105-117, (1990).
- 36. Potchter, O. et al. International Journal of Climatology, 26, 1695-1711, (2006).
- 37. Chang, C.-R. et al. Landscape and Urban Planning, 80, 386-395, (2007).
- Jo, H.-K. & McPherson, G. E. Journal of Environmental Management, 45, 109-133, (1995).
- 39. Pope, C. A. et al. Environmental Health Perspectives, 103, 472-480, (1995).
- 40. Pope, I. C. et al. JAMA, 287, 1132-1141, (2002).
- European Commission. Directive 2008/50/EC of the European Parliament and of the Council of 21 May 2008 on ambient air quality and cleaner air for Europe. (2008).
- World Health Organisation. Burden of disease from Ambient Air Pollution for 2012. (2014).

- 43. Public Health England. Estimating Local Mortality Burdens associated with Particulate Air Pollution. (2014).
- 44. Räsänen, J. V. et al. Environmental Pollution, 183, 64-70, (2013).
- 45 Freer-Smith, P. H. et al. Environmental Pollution, 133, 157-167, (2005).
- 46. Beckett, K. P. et al. Global Change Biology, 6, 995-1003, (2000).
- 47. Owen, S. M. et al. Ecological Applications, 13, 927-938, (2003).
- 48. Donovan, R. G. et al. Environmental Science & Technology, 39, 6730-6738. (2005).
- 49. Curci, G. et al. Atmospheric Environment, 43, 1444-1455, (2009).
- 50. Sartelet, K. N. et al. Atmospheric Environment, 53, 131-141, (2012).
- 51. Chameides, W. et al. Science, 241, 1473-1475, (1988).
- 52. Vos, P. E. J. et al. Environmental Pollution, 183, 113-122, (2013).
- 53. Buccolieri, R. et al. Science of The Total Environment, 407, 5247-5256, (2009).
- 54. Wania, A. et al. Journal of Environmental Management, 94, 91-101, (2012).
- 55. Pugh, T. A. M. et al. Environmental Science & Technology, 46, 7692-7699, (2012).
- 56. Nowak, D. J. et al. Urban Forestry & Urban Greening, 4, 115-123, (2006).
- 57. Nowak, D. J. et al. Atmospheric Environment, 34, 1601-1613, (2000).
- 58. Dadvand, P. et al. Science of The Total Environment, 523, 59-63, (2015).
- 59. Pauleit, S. & Duhme, F. Landscape and Urban Planning, 52, 1-20, (2000).
- Asadian, Y. & Weiler, M. Water Quality Research Journal of Canada, 44, 16-25, (2009).
- 61. Sanders, R. A. Urban Ecology, 9, 361-376, (1986).
- 62. Armson, D. et al. Urban Forestry & Urban Greening, 12, 282-286, (2013).
- 63. Bartens, J. et al. Journal of Environmental Quality, 37, 2048-2057, (2008).
- 64. Ellis, B. Applied Geography, 11, 187-200, (1991).
- Characklis, G. & Wiesner, M. Journal of Environmental Engineering, 123, 753-759, (1997).
- Jenkins, G. J. et al. UK Climate Projections: Briefing report. (Met Office Hadley Centre, Exeter, UK, 2009).
- 67. Gill, S. E. et al. Built Environment, 33, 115-133, (2007).
- 68. Villarreal, E. L. et al. Ecological Engineering, 22, 279-298, (2004).
- 69. Ellis, J. B. et al. Water and Environment Journal, 16, 286-291, (2002).
- 70. McKinney, M. L. Biological Conservation, 127, 247-260, (2006).
- 71. Cornelis, J. & Hermy, M. Landscape and Urban Planning, 69, 385-401, (2004).
- 72. Fuller, R. A. et al. Diversity and Distributions, 15, 328-337, (2009).
- 73. Schwartz, M. W. et al. BioScience, 52, 601-606, (2002).
- 74. Baldock, K. C. R. et al. Phil Trans R Soc B, 282, 20142849, (2015).
- 75. Helden, A. J. & Leather, S. R. Basic and Applied Ecology, 5, 367-377, (2004).
- 76. Saarinen, K. et al. Biological Conservation, 123, 403-412, (2005).
- 77. Brenneisen, S. Urban Habitats, 4, 27-36, (2006).
- 78. Baumann, N. Urban Habitats, 4, 37-50, (2006).
- 79. Dallimer, M. et al. BioScience, 62, 47-55, (2012).
- 80. Dallimer, M. et al. Conservation Biology, 28, 404-413, (2014).
- 81. Rouquette, J. R. et al. Diversity and Distributions, 19, 1429-1439, (2013).
- 82. Hale, J. D. et al. PLoS ONE, 7, e33300, (2012).
- 83. Potts, S. G. et al. Journal of Apicultural Research, 49, 15-22, (2010).
- 84. Goulson, D. et al. Annual Review of Entomology, 53, 191-208, (2008).
- Commission for Architecture and the Built Environment. Making contracts work for wildlife: how to encourage biodiversity in urban parks. (2006).
- Saraev, V. Economic benefits of greenspace: a critical assessment of evidence of net economic benefits. (Forestry Commission, Edinburgh, 2012).
- 87. Commission for Architecture and the Built Environment. Does money grow on trees?, (2005).
- Regeneris. The economic contribution of the Mersey Forest's Objective One-funded investments., (Regeneris Consulting, London, 2009).